



**UNIVERSITI PUTRA MALAYSIA**

**PRODUCTION AND BIOLOGICAL ACTIVITY OF MONOTERPENES  
FROM FLOWERS AND CALLUS CULTURES OF MICHELIA ALBA  
DC.**

**NOR AZIZUN RUSDI**

**FBSB 2005 7**

**PRODUCTION AND BIOLOGICAL ACTIVITY OF MONOTERPENES  
FROM FLOWERS AND CALLUS CULTURES OF *MICHELIA ALBA* DC.**

**By**

**NOR AZIZUN RUSDI**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,  
In Fulfilment of the Requirements for the Degree of Master of Science**

**November 2005**



## **DEDICATION**

With love to all my family including my mum Khatijah Said, my dad Rusdi Mat Din, my loving husband (Che Hasin Che Lah), my sisters (Nor Hakimah, Nor Nasrina, Nor Akmal, Nor Mukminah) and my brothers (Mohd Hakimi, Nasron, Muhammad and Abdul Hafiz), for enduring, with patience and understanding. Also not forgetting my lovely friends (Siti Sarah, Siti Aishah, Rozaina, Azlena, Rafidah and to my entire lab mates in lab 230) for their patience and guidance.

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Master of Science

**PRODUCTION AND BIOLOGICAL ACTIVITY OF MONOTERPENES  
FROM FLOWERS AND CALLUS CULTURES OF *MICHELIA ALBA* DC.**

**By**

**NOR AZIZUN BT RUSDI**

**NOVEMBER 2005**

**Chairman : Associate Professor Radzali Muse, PhD**

**Faculty : Biotechnology and Biomolecular Sciences**

Analysis of volatile compounds from the selected flower buds, grown *in-vitro* and callus cultures of *Michelia alba* D.C were performed using the gas chromatography technique with mass spectrometry detector (GC-MS). The results showed that the major volatile compound obtained from the three-selected flower bud was dihydrocarveol. Petal grown *in-vitro* on the Murashige and Skoog (MS) basal medium with different concentrations of plant growth regulators did not produce any dihydrocarveol. But callus grown in the culture medium containing 1.25 mg/L (w/v) of NAA plus 1.35 mg/L (w/v) of BAP (T20) and 0.8 mg/L (w/v) of NAA plus 1.85 mg/L of BAP (T23) produced linalool as a major compounds. On the other hand, callus treated with 1.25 mg/L (w/v) of NAA plus 0.45 mg/L (w/v) of BAP (T10) and T20 (1.25 mg/L (w/v) of NAA plus 1.35 mg/L (w/v) of BAP were able to synthesis cyclohexane (32.96±1.94%) compared to T23 (17.015±1.06%). Callus derived from flower petal of *M. alba* treated with 1.25 mg/L (w/v) of NAA plus 0.45 mg/L (w/v) of

BAP (T10) gave the higher growth ( $23.83 \pm 1.44$  g F.wt/ culture;  $0.23 \pm 0.09$  g D.wt/culture) respectively at pH 5.7,  $25 \pm 2^\circ\text{C}$  and in the complete dark condition after 5 weeks of culture. Treatment of *M. alba* flower petal derived callus with 1.0 mg/L (w/v) NAA plus 1.0 mg/L (w/v) BAP supplied with 150 mg/L (w/v) of casein hydrolysate showed higher growth ( $0.211 \pm 0.02$  g D.wt/ culture) than all other concentrations of casein hydrolysate. Most of the new volatile compounds in the callus cultures were found after treatment with 150 mg/L (w/v) of casein hydrolysate. Addition of various concentrations of bioelicitors, jasmonic acid (0 to 0.25 mg/mL (w/v)) yeast extract (0-0.3% (w/v)) and pectinase (0-0.25 mg/mL (w/v)) into the culture medium, decreased the cell growth. However, treatment of flower petal derived callus with (0.25 mg/mL (w/v)) of pectinase showed the highest growth ( $0.118 \pm 0.002$  g D.wt/culture) on day 18<sup>th</sup> compared to other treatment. On the other hand, petal derived callus treated with jasmonic acid and L-phenylalanine decreased the callus growth but significantly increased the production of new volatile compounds i.e. cinnamaldehyde, caryophyllene, nerolidol, cinnamic acid, ocimene, farnasene, undecene, germacrene D and also linalool. The essential oils obtained from flower bud showed the insecticidal activities, against *Tribolium castaneum* adults. The insecticidal property of essential oil obtained from flower and petal-derived callus was probably associated with presence of linalool, eugenol and geraniol. Result from seed germination assay also showed that pentane extract of *Michelia alba* flowers was able to effect the germination of the *Brassica nigra* L (mustard seeds) tested. The phytotoxicity effect and cytotoxicity action may also be possibly associated with the presence of cineol, limonene and linalool though they are minor components.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**PENGHASILAN DAN AKTIVITI BIOLOGIKAL SEBATIAN  
MONOTERPENOID DARIPADA BUNGA DAN KULTUR KALUS *MICHELLIA  
ALBA* DC.**

**Oleh**

**NOR AZIZUN BT RUSDI**

**November 2005**

**Pengerusi : Profesor Madya Radzali Muse, PhD**

**Fakulti : Bioteknologi dan Sains Biomolekul**

Kaedah kromatografi gas spektrometri jisim (GC-MS) telah digunakan untuk menganalisis kandungan sebatian meruap daripada sampel bunga cempaka putih yang dipilih. Keputusan yang diperolehi menunjukkan bahawa kandungan sebatian meruap terbanyak dari pengestrakan 3 peringkat bunga cempaka putih adalah dihidrokarveol. Kelopak bunga cempaka putih yang dikulturkan secara *in-vitro* pada medium asas pertumbuhan Murashige dan Skoog yang mengandungi pengawalatur pertumbuhan tumbuhan pada kepekatan yang berbeza gagal menghasilkan sebatian dihidrokarveol. Tetapi kultur kalus tisu yang telah ditumbuhkan pada medium yang mengandungi 1.25 mg/L (b/i) (NAA) ditambah dengan 1.35 mg/L (b/i) (BAP) (T20) dan 0.8 mg/L (b/i) (NAA) ditambah dengan 1.85 mg/L (b/i) (BAP) (T23) telah berjaya menghasilkan linalool. Selain itu, kalus yang ditumbuhkan pada medium yang

mengandung 1.25 mg/L (b/i) NAA ditambah dengan 0.45 mg/L (b/i) BAP (T10) dan T20 (1.25 mg/L (b/i) of NAA ditambah dengan 1.35 mg/L (b/i) BAP telah berjaya menghasilkan sebatian sikloheksana ( $32.96 \pm 1.94\%$ ) berbanding dengan rawatan T23 ( $17.015 \pm 1.06\%$ ). Kalus yang terhasil dari kelopak bunga cempaka putih (*M. alba*) yang diuji pada media mengandungi 1.25mg/L (b/i) (NAA) ditambah dengan 0.45mg/L (b/i) (BAP) telah memberikan pertumbuhan yang tertinggi ( $23.83 \pm 14.44$ .g berat basah/kultur;  $0.23 \pm 0.09$ g berat kering/kultur) selepas 5 minggu dieramkan pada pH 5.7, suhu  $25 \pm 2^{\circ}\text{C}$  dan dalam keadaan gelap sepenuhnya. Rawatan kalus dari kelopak bunga cempaka putih pada kepekatan 1.0 mg/L (b/i) (NAA) ditambah dengan 1.0 mg/L (b/i) (BAP) ditambah dengan 150 mg/L (b/i) kasein hidrolisat telah menunjukkan peningkatan pertumbuhan kalus ( $0.211 \pm 0.02$ g.berat kering/kultur) berbanding dengan kepekatan kasein hidrolisat yang lain. Banyak sebatian meruap yang baru telah ditemui pada kalus yang diberi rawatan dengan 150 mg/mL (b/i) kasein hidrolisat. Penambahan beberapa siri kepekatan bioelisor, asid jasmonik ( $0-0.05$ ) mg/mL (b/i), ekstrak yis ( $0-0.3\%$  (b/i)), dan pektinase ( $0-0.25$  mg/mL (b/i)) ke dalam medium asas telah menurunkan kadar pertumbuhan sel kultur tisu. Tetapi didapati rawatan kelopak bunga kalus dengan pektinase pada kepekatan 0.25 mg/mL (b/i) telah berjaya menunjukkan pertumbuhan tertinggi pada hari 18 pengamatan jika dibandingkan dengan rawatan lain. Di dalam hal lain, kelopak bunga kalus yang dirawat dengan asid jasmonik dan L-fenilalanin telah menurunkan kadar pertumbuhan kalus tetapi telah berjaya mensintesis beberapa sebatian meruap yang baru seperti sinamaldehida, karyofilin, nerolidol, asid sinamik, osimin, farnasin, undecin, germaserin D dan juga linalol. Minyak perlu yang diperolehi daripada bunga cempaka menunjukkan sifat insektisidal keatas serangga *T. castaneum* dewasa. Ciri - ciri penghindar serangga yang terdapat pada ekstrak minyak bunga cempaka putih dan

kultur kalus mungkin disebabkan oleh kehadiran beberapa sebatian seperti linalol, cineol dan limonen. Keputusan daripada asai percambahan *Brassica nigra* L (biji sawi) juga menunjukkan ekstrak bunga cempaka yang menggunakan pelarut pentana telah mempengaruhi percambahan *Brassica nigra* L (biji sawi) yang telah digunakan. Ciri-ciri fitotoksiti dan sitotoksiti yang hadir mungkin disebabkan oleh kehadiran cineol, limonen dan linalol walaupun dalam jumlah yang sedikit.



## **ACKNOWLEDGEMENTS**

I would like to sincerely thank to Assoc. Prof. Dr. Radzali Muse, my main supervisor, for his patience, guidance, and advice during this research. I also wish to thank the other members of my supervisory committee: Assoc. Prof. Dr. Johari Ramli from the Department of Biochemistry, Assoc. Prof. Dr. Md. Aspollah Sukari from the Department of Chemistry, Cik Sanimah Simoh from the Biotechnology Center (MARDI) and Encik Suri Roowi from the Food Technology Center (MARDI), for their valuable review and assistance to this research. I am deeply grateful for the opportunity to learn and work with the staff from UPM and MARDI.

I am sincerely thankful and greatly appreciative for all of the help, encouragement, guidance and friendship that I have gained during working with this project from the staff of the Department of Biochemistry, Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia; all staff of tissue culture laboratory, Biotechnology Center, Malaysian Agriculture Research and Development Institute (MARDI) Serdang. I hereby knowledge, with gratitude, Mrs Siti Shaleha, Mrs Habsah, and Ms Nor Afni Zulkifli for their help on the tissue culture techniques. I would like to give special thanks to the Pasca siswazah fellowship, UPM for two years of financial support. Finally, I would like to express my heartfelt gratitude and appreciation to my family for their advice, encouragement, understanding, support, and prayers. May God bless you all.

I certify that an Examination Committee met on 18<sup>th</sup> November 2005 to conduct the final examination of Nor Azizun Rusdi on her Master of Science thesis entitled “Production and Biological Activity of Monoterpenes from Flowers and Callus Cultures of *Michelia alba* DC” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

**Norhani Abdullah, PhD**

Professor

Faculty of Biotechnology and Biomolecular Science

Universiti Putra Malaysia

(Chairman)

**Janna Ong Abdullah, PhD**

Lecturer

Faculty of Biotechnology and Biomolecular Science

Universiti Putra Malaysia

(Internal Examiner)

**Mohd. Puad Abdullah, PhD**

Lecturer

Faculty of Biotechnology and Biomolecular Science

Universiti Putra Malaysia

(Internal Examiner)


**Shaida Fariza Sulaiman, PhD**

Associate Professor

School of Biological Sciences

Universiti Sains Malaysia

(External Examiner)



---

**HASANAH MOHD. GHAZALI, PhD**  
Professor/Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: **19 JAN 2006**

This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirements for the degree of Master of Science. Members of the Supervisory Committee are as follows:

**Radzali Muse, PhD**

Associate Professor

Faculty of Biotechnology and Biomolecular Sciences,  
Universiti Putra Malaysia.

(Chairman)

**Mohd Aspollah Hj Sukari, PhD**

Associate Professor

Faculty of Sciences,  
Universiti Putra Malaysia.

(Member)

**Johari Ramli, PhD**

Associate Professor

Faculty of Biotechnology and Biomolecular Sciences,  
Universiti Putra Malaysia.

(Member)



**AINI IDERIS, PhD**

Professor/Dean

School of Graduate Studies  
Universiti Putra Malaysia

Date: 07 FEB 2006

## DECLARATION

I hereby declare that the thesis is based on my original work except for equations and citation, which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



**NOR AZIZUN BINTI RUSDI**

Date: 16 JANUARI 2006

## TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	v
ACKNOWLEDGEMENTS	viii
APPROVAL SHEET	x
DECLARATION	xi
LIST OF TABLES	xvi
LIST OF FIGURES	xviii
LIST OF APPENDICES	xx
LIST OF ABBREVIATIONS/SYMBOLS	xxi
<b>CHAPTER</b>	
<b>I</b> INTRODUCTION	1
Objectives of The Study	3
<b>II</b> LITERATURE REVIEW	
<i>Michelia alba</i> or ‘Cempaka Putih’.	4
Plant Cell and Tissue Culture as Source of Secondary Metabolites.	5
<i>M. alba</i> (Cempaka Putih) Flowers.	8
Techniques in Samples Preparation and Identification of Chemical Composition of Monoterpenes	9
Extraction	10
Solvent Extraction	11
Simultaneous Distillation Adsorption	12
Simultaneous Distillation Extraction	14
Steam Distillation Extraction	18
Gas Chromatography	18
Composition of Essential Oils	19
Production of Essential Oils	20
Location of Essential Oils in Plants	20
Problems in Producing Essential Oils in Plant Cell and Tissue Culture	22
Manipulation of Nutrient to Improve Yield	24
Physical Factors	26
Effect of different Incubation (light-dark) on Callus growth of <i>M. alba</i>	27
Effect of different Initial pH Medium on callus growth of <i>M. alba</i>	27
Chemical Factors	28
Effect of Plant Growth Regulators on Callus Growth of <i>M. alba</i>	28



	Effect of different Carbon Sources on Callus Growth of <i>M. alba</i>	32
	Stress in Plant Tissue Culture	34
	Effect of Precursors on Callus Growth for Optimum Production of Monoterpene Compounds	35
	Effect of Elicitors on Callus Growth for Optimum Production of Monoterpene Compounds	36
	Biological Activities of Essential Oils (Monoterpenes)	38
	Allelopathic Activity	39
	Insecticidal Activity	39
<b>III</b>	<b>MATERIALS AND METHODS</b>	
	Common Chemicals	42
	Preparation of Stock Solution of Plant Growth Regulators (PGRs)	42
	Preparation of Plant Tissue Culture Medium	43
	Plant Materials	43
	<i>Michelia alba</i> Explants	46
	Initiation of <i>M. alba</i> callus	46
	Growth of <i>M. alba</i> tissue cultures	47
	Determination of Callus Growth for Optimum Production of Monoterpene Compounds	48
	Effect of Different concentrations of casein hydrolysate (CH) on callus growth	49
	Phytohormone Treatments on callus growth of <i>M. alba</i>	49
	Effect of different incubation (light-dark) on callus growth of <i>M. alba</i>	50
	Effect of Different initial pH on callus growth of <i>M. alba</i>	50
	Effect of Different concentration of Carbon Sources on callus growth of <i>M. alba</i>	50
	Effect of different concentrations of sucrose on callus growth of <i>M. alba</i>	51
	Effect of Different Type of Elicitors and Precursors on callus growth of <i>M. alba</i>	51
	Determination of Essential Oil from callus and Intact Plant of <i>Michelia alba</i> flowers	
	Isolation of Essential oil by Simultaneous Distillation Extraction (SDE) Apparatus	52
	Gas Chromatography Condition	54
	Biological Activities of <i>M. alba</i> Essential Oils	54
	Germination and Growth Test	55
	Insecticidal Activities of Monoterpenes	
	Preparation of Artificial Diet Media (ADM)	56
	<i>Tribolium castaneum</i> Adults Feeding	56
	Insect Repellent Test	56
	Standard Chemicals Used	57
	Contact Toxicity	58
	Statistical Analysis	58

<b>IV</b>	<b>RESULTS AND DISCUSSIONS</b>	
	<b>Production of Essential Oils (Monoterpenes Compounds)</b>	<b>60</b>
	<b>Effect of Explants Materials</b>	
	<b>Volatiles Compounds Present in different Parts of</b>	<b>64</b>
	<i>M. alba</i> Plants	
	<b>Plant Tissue cultures</b>	
	<b>Application of Plant Growth Regulators</b>	<b>65</b>
	<b>Effect of different Concentration of NAA and BAP on</b>	<b>67</b>
	initiation of <i>M. alba</i> callus Derived from Petal	
	<b>PGRs influence of IBA, ZEATIN, 2, 4-D, NAA and</b>	<b>69</b>
	BAP on initiation of the <i>M. alba</i> callus Derived from	
	Petal	
	<b>Effect of different concentration of BAP and 2,4-D on</b>	<b>70</b>
	initiation of <i>M. alba</i> callus Derived from Petal	
	<b>Polivinylypyrrolidone (PVP) Supplementation</b>	<b>72</b>
	<b>Effect of different concentration of Casein Hydrolysate</b>	<b>74</b>
	on callus Growth of <i>M. alba</i> .	
	<b>Effect of different concentration of Casein Hydrolysate</b>	<b>76</b>
	on Production of monoterpenes	
	<b>Effect of NAA and BAP Supplied on Yellow Bud</b>	<b>79</b>
	Derived Callus Growth	
	<b>Effect of NAA and BAP Supplied on Production of</b>	<b>82</b>
	Volatile Oil Compounds (monoterpenes)	
	<b>Effect of Different Sucrose Concentrations on Callus</b>	<b>84</b>
	Growth of <i>M. alba</i>	
	<b>Effect of Carbon Sources on Callus Growth of <i>M. alba</i></b>	<b>87</b>
	<b>Effect of different Initial pH Medium on Callus Growth</b>	<b>90</b>
	of <i>M. alba</i>	
	<b>Effect of different Incubation (Light- Dark) on Callus</b>	<b>93</b>
	Growth of <i>M. alba</i>	
	<b>Effect of Light Incubation on Production of Volatile</b>	<b>95</b>
	Oil Compounds	
	<b>Effect of Different Types of Agar on <i>M. alba</i> Callus</b>	<b>96</b>
	Cultures	
	<b>Production of Monoterpenes Compounds in callus Derived</b>	<b>99</b>
	<b>from Different Stages of Flower Buds.</b>	
	<b>Effect of Elicitors</b>	<b>99</b>
	<b>Effect of Elicitors on Yellow Bud Derived Callus Growth</b>	<b>100</b>
	<b>(After 3 weeks)</b>	
	<b>Effect of Jasmonic acid, Pectinase and Yeast Extract on the</b>	<b>102</b>
	<b>Production of Volatile Oil Compounds</b>	
	<b>Effect of L-Phenylalanine on the Production of Volatile Oil</b>	<b>105</b>
	<b>Compounds</b>	

Effect of L-Phenylalanine on Petal Derived Callus Growth	107
Biological Activities of <i>M. alba</i> Essential Oils	108
Allelopathic Activity of <i>M. alba</i> extracts	109
Germination Percentage	109
Radicle Length	112
Seedling Weight	112
Insecticide Activity	115
Insect Repellency Testing	116
Contact Toxicity	118
<b>V</b> <b>CONCLUSION</b>	<b>122</b>
<b>REFERENCES</b>	<b>146</b>
<b>APPENDICES</b>	<b>147</b>
<b>BIODATA OF THE AUTHOR</b>	<b>153</b>



## LIST OF TABLES

Table		Page
1	List of group of natural products that were so far isolated from tissue and suspension cultures of higher plants.....	6
2	Methods for isolation and concentration of volatiles oil.....	10
3	The retention time and all the volatile compounds found in the three types of flower buds of <i>M. alba</i> (S5, S6 and S7).....	62
4	Type of full strength of basal MS medium supplemented with various combinations of NAA (mg/mL) plus BAP (mg/mL).....	60
5	Type of half strength of basal MS medium supplemented with various combination s of 2,4-D mg/mL plus BAP mg/mL.....	61
4	The major volatile compounds present in three parts (Leaf, Stem and Root) of <i>M. alba</i> Plant.....	82
5	Callus initiation of <i>M. alba</i> from petal explants in the present of PGRs: naphthalene acetic acid (NAA), benzylaminopurine (BAP), indole acetic acid (IAA), kinetin (KIN), 2,4-dichlorophenoxy acetic acid (2,4-D).....	66
6	Callus initiation of <i>M. alba</i> from petal explants in the present of PGRs: naphthalene acetic acid (NAA) plus benzylaminopurine (BAP) .....	68
7	Callus initiation of <i>M. alba</i> from petal explants in the present of PGRs : naphthalene acetic acid (NAA), benzylaminopurine (BAP), Zeatin (ZEA), indole-butyric acid (IBA).....	70
8	Callus initiation of callus of <i>M. alba</i> bud explants in the presence of 2,4-dichlorophenoxy acetic acid (2,4-D) and benzylaminopurine (BAP) in Half Strength of MS Basal Medium.....	72
9	Chemical compounds in essential oil of four cell lines <i>M. alba</i> callus cultures and their relative percentage in the oils.....	78
10	Volatile compounds present in callus derived from petal of <i>M. alba</i> grown at 25±2°C and on Light Condition.....	76

11	Chemical compounds present in the <i>M. alba</i> callus cultures treated with elicitors (pectinase and jasmonic acid) and their relative percentage in the oils.....	104
12	Effect of <i>M. alba</i> flowers extract and three volatile monoterpenes on the radicle length, seedling weight of seedling of mustard seed ( <i>Brassica nigra</i> L) after 168 h exposure time.....	114
13	Effect of dilute extracts of <i>M. alba</i> flowers and three volatile monoterpenes on percentage of germination and seed vigour of mustard seed ( <i>Brassica nigra</i> L) after a week exposure time at each 8 mg/mL of concentration.....	115
14	Contact toxicity of six types of standard volatile monoterpenes (Linalool, eugenol, cineol, $\alpha$ -pinene and limonene) against <i>T. castaneum</i> adults after 96 h exposure time.....	121

## LIST OF FIGURES

Figure		Page
1	A Simultaneous Distillation Adsorption (SDA).....	13
2	A Modified Lickens and Nickerson Apparatus (SDE).....	17
3	A Schematic Representation of Gas Chromatograph.....	19
4	A Schematic Diagram of production under stress. Various stress factors affecting the production of secondary metabolites.....	34
5	Various stages of flower development of <i>M. alba</i> (Cempaka Putih).....	45
6	Callus induced from petals segment of <i>M. alba</i> flower on a basal MS medium (pH 5.7) supplied with 250mg/L (w/v) of casein hydrolysate after four weeks of culturing .....	47
7	Modified Lickens and Nickerson Apparatus of Simultaneous Distillation Extraction (SDE) for Essential oil Extraction.....	53
8	Initiation and Use of callus and cell suspension cultures of <i>Michelia alba</i> .....	59
9	The level of dihydrocarveol; butanoic acid, acid 2-methyl, methyl ester; cyclohexane, 1-ethenyl-1-methyl-2, 4-bis (1-methylethenyl); eugenol methyl ether and bicyclo[7.2.0] undec-4-ene, 4, 11,11 trimethyl-8-methylene throughout stages of flower bud formation on S5, S6 & S7.....	64
10	Percentage of growth for <i>M. alba</i> callus derived from petal explants on basal MS medium supplied with 0.5 and 1.0% (w/v) of polyvinylpyrrolidone (PVP) .....	73
11	Fresh weight (a) and Dry weight (b) of <i>M. alba</i> callus culture grown on a MS basal medium supplied with four concentrations of casein hydrolysate for 5 weeks incubation.....	75
12	Fresh weight (a) and dry weight (b) of <i>M. alba</i> callus culture grown on four solid basal media (MS) supplied with different concentrations of NAA plus BAP after 5 weeks incubation.....	80
13	Effect of supplied NAA plus BAP on <i>M. alba</i> petals-derived callus growth after 4 weeks of culturing.....	81

14	Effect of supplied NAA plus BAP on the production of volatile oil, compounds monoterpenes in <i>M. alba</i> petal derived callus.....	83
15	Time course of dry weight of <i>M. alba</i> callus culture in different concentrations of sucrose such as 2, 3 and 4% (w/v) and casein hydrolysate supplied such as (a) 25; (b) 50; (c) 100; (d) 150 mg/L (w/v) of CH .....	86
16	Time course of dry weight of <i>M. alba</i> callus cultures of four- cell lines in different carbon sources and casein hydrolysate) (CH) supplied such as (a) 25; (b) 50; (c) 100; (d) 150 mg/L (w/v) of CH.....	89
17	Time course of dry weight of <i>M. alba</i> callus cultures of four- cell lines in different pH values.....	92
18	The effects of light and dark condition on growth of <i>M. alba</i> callus cultures containing four cell lines such as (a) 25; (b) 50; (c) 100; (d) 150 mg/L (w/v) of casein hydrolysate (CH).....	94
19	The effect of two types of agar (gelrite and kelcogel) used in a basal MS medium on the growth of <i>M. alba</i> callus cultures supplied (a) 25; (b) 50; (c) 100; (d) 150 mg/L (w/v) of casein hydrolysate (CH).....	98
20	Effect of different elicitors used on biomass accumulation in callus cultures of <i>M. alba</i> .....	101
21	Effect of supplied elicitors (Jasmonic acid and Pectinase) on the production of volatile oil compounds in <i>M. alba</i> petal derived callus.....	102
22	Effect of supplied L-phenylalanine on the production of volatile Oil compounds in <i>M. alba</i> petal-derived callus cultures.....	106
23	Effect of different concentrations of L-Phenylalanine used on biomass accumulation in callus cultures of <i>M. alba</i> .....	108
24	Effect of n-pentane extract of <i>M. alba</i> flowers at concentration 8 mg/mL on growth of the <i>Brassica nigra</i> L .....	110

25	Effect of two volatile monoterpenes at concentration 8mg/mL on the growth of <i>Brassica nigra</i> L.....	111
26	Number of <i>Tribolium castaneum</i> adults on petri dish with artificial diet media (ADM) after 4 weeks incubation (left); <i>T. castaneum</i> adult under microscope with magnification, $M=\times 10$	115
27	Amount of <i>T. castaneum</i> adults left in artificial diet medium (ADM) after 24 h exposure time, at $27\pm 2^{\circ}\text{C}$ .....	117
28	Effect of <i>M. alba</i> essential oils on the mortality of <i>T. castaneum</i> adults after 96 h exposure time at $27\pm 2^{\circ}\text{C}$ .....	119
29	Mortality of <i>T. castaneum</i> caused by different concentrations of <i>M. alba</i> flowers oil and six (6) types of standard volatile monoterpenes (linalool, geraniol, eugenol, cineol, limonene and $\alpha$ -pinene) after 96 h exposure time at $27\pm 2^{\circ}\text{C}$ .....	120

## LIST OF APPENDICES

Appendix	Page
A      Composition of Murashige and Skoog (MS) (1962) Basal Medium.....	147
B (a)    Type of full strength basal MS medium supplemented with various combinations of NAA mg/L (w/v) plus BAP mg/L (w/v).....	148
B (b)    Type of half strength basal medium supplemented with various combinations of 2, 4-D mg/L (w/v) plus BAP mg/L (w/v).....	148
C-1 (a) <i>M. alba</i> stamen and ovary cultured on the basal half strength of MS medium containing phytohormones (0.06 mg/L (w/v) of 2, 4-D and 0.1 mg/L (w/v) of BAP.....	149
C-1 (b) <i>M. alba</i> stamen cultured on the basal half strength of MS medium containing phytohormone (0.03 mg/L (w/v) of 2, 4-D and 0.2 mg/L (w/v) of BAP) at pH 5.7, 25±2°C, after 3 months of culturing in the complete dark condition.....	149
C-2      Callus cultured after 1-month inoculation on a basal MS medium containing phytohormone (1.0 mg/L (w/v) of NAA plus 1.0 mg/L (w/v) of BAP).....	150
D-1      GC-MS Chromatogram of extracted Volatile Compounds in Callus Derived from (a) treatment with 0.8 mg/L (w/v) NAA plus 1.85 mg/L (w/v) BAP, (b) treatment with 1.25 mg/L (w/v) NAA plus 1.35 mg/L (w/v) BAP.....	151
D-2      GC-MS chromatogram of Volatile oil Compounds extracted from three (3) types of <i>M. alba</i> flower buds. (a) Stage 5; (b) stage 6; (c) stage 7. ....	152

## LIST OF ABBREVIATIONS AND SYMBOLS

Abs	Absorbance
ADM	Artificial diet media
App	Appendix
Approx.	Approximately or about
Aq	Aqueous
BAP	6-benzylaminopurine
b/i	berat per isipadu
b.p	Boiling point
°C	Degrees Celsius
CF(s)	Culture filtrate (s)
CH	Casein Hydrolysate
Conc. (s)	Concentrations
2, 4-D	2, 4-dichlorophenoxy acetic acid
d	Day
D.wt	Dry weight
e.g.	Example (s)
<i>et al.</i> ,	<i>at alli</i> and other people
etc.	<i>et cetra</i> , and the rest.
EB	Evans blue (stain)
EO	Essential oil
EtOH	Ethanol or ethyl alcohol
FeNaEDTA	Ethylenediaminetetraacetic acid, Ferric Sodium salt trihydrate,
F.wt	Fresh weight
GC	Gas Chromatography
GC-MS	Gas chromatography mass spectrometry
h	Hour
IBA	Indole butyric acid
IAA	Indole acetic acid
JA	Jasmonic acid
KIN	Kinetin
LD <sub>50</sub>	Lethal dose at 50%
LD <sub>90</sub>	Lethal dose at 90%
MS	Murashige and Skoog
NAA	Naphthalene Acetic Acid
PVP	Polyvinylpyrrolidone
SD	Standard deviation
SDE	Simultaneous Distillation Extraction
SEM	Standard error of mean
μl	Micro liter
w/v	weight per volume
ZEA	Zeatin

## CHAPTER I

### INTRODUCTION

For centuries, mankind is totally dependent on plants as sources of carbohydrates, proteins and fats for food shelter. In addition, plants are valuable sources of a wide range of secondary metabolites, which are used as biopharmaceuticals, agrochemicals, flavours, fragrances, colours, biopesticides and food additives as reported by Balandrin and Klocke (1988).

Plants produce a variety of chemical compounds that have large economical value. Harborne (1991) reported that first of all the chemical compounds are connected with important traits of the plant itself, *e.g.*, colour or fragrance of flowers, taste and color of foods, and resistance against pests and diseases and also for the production of fine chemicals such as drugs, antioxidants, flavours, fragrances, dyes, insecticides and pheromones.

Monoterpenes such as linalool, champhene and cineol, are involved in pollinator attraction and allelopathy (Langenheim, 1994; Harborne, 1991). Monoterpenes has also found expensive industrial applications as flavoring agents, topical medicines, perfumes and insecticides (Croteau *et al.*, 2000; Little and Croteau, 1999; Loza-Tavera, 1999; Phillips and Croteau, 1999).



Monoterpenes are widespread in the plant kingdom (Banthorpe and Charlwood, 1980) and are often responsible for the characteristic odours of plants. These substances are believed to function principally in ecological roles, serving as herbivore-feeding deterrents, antifungal defense, and attractants for pollinators (Langenheim, 1994; Romagni *et al.*, 2000). There is also evidence that monoterpenes have potential uses as herbicides, pesticides, antimicrobial agents and dietary anticarcinogens (Crowell, 1999). The monoterpenoids are the major component of many essential oils and as such, have economic importance as flavors and perfumes (Briellmann, 1998).

*Michelia alba* flowers is a native plant of tropical and subtropical Southeast Asia. *Michelia* ('cempaka') species is interestingly known to produce high value essential oil. The popularity of the fragrance has resulted in many species of *Michelia* being grown all over the world. It is reported that the essential oil of *Michelia* has been used as a key ingredient in Joy and J'adore, among the world's most expensive perfumes. In China, it is reported by Yamanishi *et al.* (1988) that the flowers are used to scent Yulan tea. The chemical composition of a few species of *Michelia* spp has been the subject of limited study.

Many plant secondary metabolites produced by higher plants are of economically importance (Stafford *et al.*, 1986; Parr, 1989). Many secondary metabolites are obtained directly from tissues whose availability may be limited. Plant cell culture offer an alternative for the production of these metabolites because the cells can be cultured in large quantities under controlled conditions and harvesting of the desired product can be less problematic (Parr, 1989; Buitelaar and Trampller, 1992).